

Review Article

A Systematic Review on Effects of Nitrogen Fertilizer Levels on Cabbage (*Brassica oleracea* var. *capitata* L.) Production in Ethiopia

Yohannes Gelaye 

Department of Horticulture, College of Agriculture and Natural Resources, Debre Markos University,
P.O. Box 269, Debre Markos, Ethiopia

Correspondence should be addressed to Yohannes Gelaye; yohanes_gelaye@dmu.edu.et

Received 22 February 2024; Revised 26 March 2024; Accepted 18 April 2024; Published 30 April 2024

Academic Editor: Abreham Berta Aneseyee

Copyright © 2024 Yohannes Gelaye. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cabbage (*Brassica oleracea* var. *capitata* L.) holds significant agricultural and nutritional importance in Ethiopia; yet, its production faces challenges, including suboptimal nitrogen fertilizer management. The aim of this review was to review the possible effect of nitrogen fertilizer levels on the production of cabbage in Ethiopia. Nitrogen fertilization significantly influences cabbage yield and quality. Moderate to high levels of nitrogen application enhance plant growth, leaf area, head weight, and yield. However, excessive nitrogen levels can lead to adverse effects such as delayed maturity, increased susceptibility to pests and diseases, and reduced postharvest quality. In Ethiopia, small-scale farmers use different nitrogen levels for cabbage cultivation. In Ethiopia, NPSB or NPSBZN fertilizers are widely employed for the growing of various crops such as cabbage. 242 kg of NPS and 79 kg of urea are the blanket recommendation for the current production of cabbage in Ethiopia. The existing rate is not conducive for farmers. Therefore, small-scale farmers ought to utilize an optimal and cost-effective nitrogen rate to boost the cabbage yield. Furthermore, the effectiveness of nitrogen fertilization is influenced by various factors including the soil type, climate, cabbage variety, and agronomic practices. Integrated nutrient management approaches, combining nitrogen fertilizers with organic amendments or other nutrients, have shown promise in optimizing cabbage production while minimizing environmental impacts. The government ought to heed suggestions concerning soil characteristics such as the soil type, fertility, and additional factors such as the soil pH level and soil moisture contents.

1. Introduction

Cabbage, a member of the *Cruciferae* (*Brassicaceae*) family, is a biennial plant often cultivated annually, characterized by a short stem and dense cluster of leaves forming a compact head [1]. It has been reported that they are closely related to broccoli, cauliflower, and Brussels sprouts and originate from Western Europe and the Northern shores of the Mediterranean [2]. Cabbage is grown in more than 90 countries worldwide [3]. China and India are typically the top producers, followed by South Korea, Germany, Japan, and South Africa [4]. South Africa, Kenya, Egypt, Ethiopia, and Niger are the top five cabbage producers on the African continent. It ranks fifth among vegetable crops worldwide [5]. Cabbage, a popular cool season crop among gardeners

and commercial producers, stands as one of the most significant vegetable crops cultivated worldwide [6]. It has been domesticated since ancient times and is renowned for its nutritional value and abundant mineral and vitamin content, including vitamins A, B1, B2, and C, making it a staple in human consumption [7]. Cabbage is also known for its cooling effect as an appetizer that can help prevent constipation [8] and protect against cancer [9]. Although cabbage prefers cool and moist climates, it can grow easily under a wide range of environmental conditions in both temperate and tropical regions [10]. Cabbage, alongside chili, holds the distinction of being Ethiopia's second most vital vegetable crop, both in terms of widespread cultivation and production [5]. However, owing to the scarcity and high cost of chemical fertilizers, most small-scale farmers in the

country employ fertilizer levels lower than the recommended rate [11]. While the ideal yield is determined by applying 79 kg/ha, there has not been enough research into how varying rates of N-fertilizer application affect the yield and yield components of cabbage [12]. Hence, the yield obtained was reported to be low (6–20 tons/ha) [13]. Similarly, declining produce quality and soil fertility in Ethiopia pose significant challenges for market acceptance and production. Nitrogen's role in cabbage yield varies with the soil type, soil fertility, and environmental factors. Research indicates that proper management of nitrogen application in terms of rate and timing can boost the cabbage yield [14]. To secure high cabbage yield, it is also important to use the optimum rate of nitrogen per unit area [15]. The ideal nitrogen rate crucially affects solar radiation absorption in cabbage crops. Despite global research, Ethiopia lacks specific studies on this aspect, which are impacted by local factors such as soil and climate. Understanding how varieties and conditions dictate optimal nitrogen levels is vital for sustainable cabbage farming, enabling the balancing of increased production with environmental concerns through tailored management practices [16]. Continuous high nitrogen use can harm soil health, necessitating long-term studies, while climate change necessitates resilient fertilizer strategies, adding complexity [17]. To benefit Ethiopian farmers, research must bridge the gap between findings and practical application through knowledge-sharing initiatives and extension services. Thus, addressing these gaps can enhance our understanding of nitrogen fertilizer's impact on cabbage production in Ethiopia, bolstering sustainable agricultural development in the region. Therefore, the aim of this review was to review the effect of nitrogen fertilizer levels on the growth and yield components of cabbage.

2. Review Methodology

In this review of the literature, I employed diverse approaches. I referred to respected journals sourced from Scopus, Web of Science, and PubMed databases. The selection criteria predominantly targeted articles released post-2019, while omitting pertinent data and books (Figure 1). Therefore, the review materials were chosen spanning the years, guided by indices and respective protocols. In total, 32 articles (30.4%), 37 articles (35%), 34 articles (32%), 25 articles (23%), and 37 articles (35%) were sourced from WoS, Scopus, PubMed, All, and other databases, respectively. The year-wise analysis indicates that the majority of articles were published between 2019 and 2023 (Figure 1). The majority of the reviewed papers concluded that varying levels of nitrogen fertilizer significantly impact the growth, yield, and quality components of cabbage crops.

3. Origin and Distribution of Cabbage

Cabbage (*Brassica oleracea* var. *capitata*) is believed to have originated from the Mediterranean region, specifically from the wild cabbage plant native to coastal areas of Europe [3]. It has a long history of cultivation, dating back thousands of years [18]. In Ethiopia, cabbage is thought to have been

introduced during the colonial period, primarily by European settlers and missionaries [19]. Cabbage cultivation in Ethiopia has grown, mainly in highland regions due to favorable climate [20]. These regions encompass the central highlands, including Addis Ababa, and the northern highlands, home to cities such as Bahir Dar and Gondar. Cabbage cultivation in Ethiopia has increased significantly in recent years due to its adaptability to various agroecological zones, its nutritional value, and its economic importance [21]. It is cultivated both by small-scale farmers for local consumption and by larger commercial farms for export markets [22]. Overall, cabbage significantly contributes to Ethiopian agriculture and diet, enhancing food security and farmer income nationwide.

4. Description of Cabbage

Cabbage has a chromosome number of $2n = 2x = 18$ and has been grown for more 3000 years in the Mediterranean region [23]. It is described as originating in Western Europe, with temperate climates, and a kale-like ancestor was grown in gardens as far back as the time of the Roman Empire [24]. Cabbage is described as a biennial crop although it is grown as an annual crop, and there is a great difference between the cultivated types of cabbage [25]. It is also known to have an adventitious root system and an unbranched stem that remains less than 30 cm long [26]. As cabbage grows, leaves multiply to form a ball-shaped "head" at the plant's centre, primarily composed of a large vegetative terminal bud resulting from a leaf overlying its shortened stem [27]. An example from a 2010 Ethiopian guide on cool season vegetable crops, seed production, and development practices describes head cabbage (common cabbage) as having a dense head of leaves, a short stem, and additional edible leaves [28]. The diverse shapes and sizes of cabbage heads across varieties encompass oblong, oval, or nearly circular forms, with cultivars displaying distinctions in the leaf size, shape, color, and head texture [29]. Clear descriptions of cabbage growth stages, though lacking standard terminology, are vital for pest management, as plant susceptibility varies, as outlined in subsequent literature (Table 1).

5. Cabbage Production in Ethiopia

The annual production of cabbage in the world is estimated to be about 70,644,191 tons with an average productivity of about 29.23 tons/ha [30]. The report from the Central Statistical Agency (CSA) in Ethiopia details 5150.14 hectares of land, yielding 35,123.65 tons of produce with a productivity rate of 6.8 tons per hectare, falling below the global average [31]. Cabbage is frequently produced during the rainy season although some commercial farmers produce it during the dry season using irrigation [32]. In Ethiopia, cabbage is cultivated both in mixed cropping systems and monoculture, with irrigation noted to significantly increase cabbage yield compared to rain-fed conditions. Cabbage cultivation is scattered throughout the highlands of Ethiopia. Cabbage production in Ethiopia centers on Copenhagen and early




FIGURE 1: Graphical representations depicting tactics for gathering articles related to cabbage cultivation.

TABLE 1: Growth stages of cabbage.

S/n	Growth stages	Descriptions	References
1	Stage 1: cotyledons (Seed leaves)	No true leaves present	[103–105]
2	Stage 2: seedling	Up to 5 true leaves	
3	Stage 3: 6–8 true leaves	Ready for transplanting	
4	Stage 4: 9–12 true leaves	Base of stem still visible from above	
5	Stage 5: precupping	Approximately 13–19 leaves	
6	Stage 6: cupping	Approximately 20–26 leaves	
7	Stage 7: early head formation	Head diameter will be approximately 10 cm	
8	Stage 8: head fill	Head diameter will be approximately 10–20 cm, and a firm round head is visible within the wrapper leaves	
9	Stage 9: mature	Head diameter will be approximately 15–30 cm, and no new visible leaf production will occur after the head has attained maximum hardness and size. The head is reported to be ready for harvest and may split if not harvested in time	

drum head varieties in the central highlands, favored by urban landholders and characterized by shallow depth, contrasting with rural farmers.

6. Importance of Cabbage

Cabbage is an economically and nutritionally important vegetable crop worldwide and is one of the most extensively consumed vegetable crops in Ethiopia, particularly in urban areas [33]. It has been reported as one of the top 20 vegetables used as a global food source [34]. Cabbage, prized for its leaves, is a rich source of minerals such as calcium, iron, sodium, potassium, and phosphorus, as well as essential vitamins, proteins, and carbohydrates [6]. Head cabbages have significant quantities of nutrients and for instance, a 100 g edible portion of cabbage is reported to contain 27 mg of calories, 1.8 mg protein, 0.1 mg fat, 4.6 mg

carbohydrate, 0.6 mg mineral, 29 mg calcium, 0.8 mg iron, and 14.1 mg sodium (Figure 2). Cabbage, rich in beta-carotene and ascorbic acid, also contributes to a healthy diet with glutamine, amino acids, and dietary glucosinolates, supporting human health [35]. Similarly, it has also been employed in the preparation of soup mixtures with other vegetables, such as carrots, celery, corn, and onions [36]. Currently, consumers in Ethiopia are aware of about cabbage crops, which are becoming more prevalent because of their nutritional value and anticancer properties.

7. Ecological Requirements of Cabbage

Cabbage is grown within an altitude range of 700–2,200 m.a.s.l [37]. In low-altitude regions, crop cultivation, particularly of water-intensive crops like cabbage, should be timed for the cooler months of the year to align




FIGURE 2: Nutritional composition of cabbage [6].

with Ethiopia's specified rainfall requirement of 380–500 mm, evenly distributed throughout the growing period [10]. Cabbage's water needs vary with soil, growth stage, and environmental factors, thriving in cooler temperatures with an optimal production range of 16–20°C (Figure 3) [38]. Temperatures exceeding 25°C have been observed to impede head formation in cabbage, which thrives in well-drained sandy or silty loam with a high organic matter content [39]. Correspondingly, the optimal soil pH range for cabbage cultivation is 6.0–6.5 [40]. Cabbage, like most leafy vegetables, is considered a heavy feeder and does well in soils with high nitrogen content [41]. Nitrogen, phosphorus, and sodium are vital for cabbage growth, with nitrogen specifically noted for fostering rapid growth, high yield, and quality [42]. In addition, the level of nitrogen is reported to lead to a higher dry matter content in the shoots than in the roots [43]. Experts conducting research identified a strong correlation between nitrogen fertilizer application and cabbage quality, recommending 242 kg of NPS and 79 kg of urea for cabbage cultivation in Ethiopia [44]. Cabbage is harmful to light frost and is attacked by many pests and diseases (Figure 3) [45]. Typically, cabbage is transplanted and shows a good response to nitrogen [46].

8. The Effect of Nitrogen on the Cabbage Yield

Cabbage growers in Brazil use blanket fertilizer recommendations, including nitrogen, at different rates and times of application [47]. Cabbage is a highly demanding vegetable crop for major plant nutrients, including nitrogen. The doubling of agricultural food production worldwide over the past four decades has been linked to a sevenfold increase in the use of nitrogen fertilizers [48]. The ongoing discovery of nitrogen fertilizers in agriculture will significantly impact the industry, exemplified by cabbage absorbing nitrogen from

organic matter such as nitrate or ammonium [49]. Nitrogen is reported to play a role in protein formation as a component of chlorophyll; thus, an adequate supply of nitrogen increases the efficiency of chlorophyll [50]. In Bangladesh, from 2013 to 2014, a study assessed the impact of varying nitrogen doses on cabbage growth and yield, suggesting that utilizing 200 kg N/ha could be beneficial for cabbage cultivation [51]. If a plant is supplied with an optimum amount of nitrogen, there is a tendency to increase leaf cell number and cell size, with an overall increase in leaf production. Moreover, the cabbage cultivation manual indicates that cabbage requires plenty of NPK nutrients for head formation [52]. In Pakistan, application of NPK farmyard manure (FYM) at a ratio of 160 : 90 : 60 along with 15–20 tons/ha of farmyard manure is reported to have a desirable effect on the growth and marketable yield of cabbage [53]. A study conducted to evaluate the impact of increasing plant populations combined with different doses of nitrogen on the yield of cabbage, its structure, and plant composition indicated that the enhancement of N dose to 300 kg N/ha was beneficial for the total yield of heads as well as those weighted greater than 1.0 kg [54]. The sole alteration in plant composition due to heavy nitrogen fertilization was the rise in nitrate accumulation and reduction in calcium content in cabbage heads, emphasizing that optimal nitrogen and nutrient levels ensure a well-balanced mineral nutrition profile [55]. Imbalanced nitrogen application harms cabbage quality, causing loose-head formation and decay; however, employing techniques such as high-density planting and careful fertilization can help control the size. A field experiment conducted to evaluate the potential of using bioslurry as organic manure for cabbage production in Bangladesh stated that it is possible to reduce 30% of the recommended nitrogen and K fertilizers and 100% of the phosphorus and sulfur fertilizers through the use of poultry or cow dung bioslurry in cabbage production [56, 57].




FIGURE 3: Ecological necessities of cabbage crop.

Research on Chinese cabbage in organic farming showed significant impacts on growth, yield, and quality by combining bioslurry, derived from anaerobic decomposition of organic materials, with inorganic fertilizers, thereby sustaining crop production and enhancing soil health [58, 59]. Manure application can reduce invalid crop water consumption, use soil water reasonably, and improve water use efficiency at different growth stages of a crop.

An experiment on head cabbage yield and quality found that decreased N and K residuals postharvest suggested complete fertilizer absorption, which is crucial ecologically [60]. Organic manure and biofertilizers improve soil health and crop quality, as evidenced by a study on red cabbage cultivation assessing nutrient efficiency, productivity, and soil sustainability [15]. In addition, a field experiment was conducted to evaluate the growth, productivity, and economics of cabbage in the grid zone as influenced by different levels of zinc and sulfur reported that growth parameters such as plant height, plant spread, number of leaves per plant at all crop growth stages, yield attributing characters, length and diameter of head and weight per head, cabbage yield per hectare net return, and benefit cost ratio increased with increasing levels of sulfur from 20 to 80 kg/ha and zinc from 2 to 8 kg/ha and the crop responded only up to 60 kg S/ha and 6 kg Zn/ha [61, 62]. During a 2017 experiment in Ghana's Kintampo North Municipality, evaluating cabbage response to various soil amendments from August to November, poultry manure matched NPK (15-15-15) and outperformed other treatments across all measured parameters, while cow dung and goat manure also exhibited promising results [63, 64]. Likewise, a field experiment conducted to determine the optimum combination of FYM and biofertilizers for improving the growth, yield, and quality attributes of Chinese cabbage under high-altitude

rain-fed climatic conditions reported that among the treatments studied, the highest plant height (33.33 cm), number of wrapper leaves/plant (18.00), plant spread/plant (111.66 cm), head weight/plant (553.33 g), yield/plot (15.49 kg), yield/ha 34.42 tons, moisture content 98.00%, and vitamin C 28 mg/100 g were observed with the application of FYM at 25 tons/ha along with an azotobacter and phosphate-solubilizing bacteria [65].

An experiment was conducted to evaluate the performance of three cabbage varieties (*Brassica oleraceae* var. *capitata* L.) varieties (Benelli, "Cairo" and "Paradox") (Bahamas-2012) indicated that the variety "Paradox" produced the longest head length and the largest head width [66]. The variety produced the highest head weight and best yield potential of 37.7 tons/ha, while combining vermicompost with integrated plant nutrients and recommended chemical fertilizers in a field experiment resulted in superior cabbage growth and yield compared to using chemical fertilizers alone [67].

During field trials conducted at the Umsunduze Training Centre in KwaZulu Natal in 2005 and 2006, exploring the effects of three organic fertilizers (chicken, Kraal manure, and compost) on cabbage (Conquistador cultivar) and carrots (Kuroda cultivar) growth, yield, and quality, it was found that cabbage head diameter was consistent between those treated with chicken manure and inorganic fertilizer [68]. A study on cabbage yield and head weight revealed that, on average, heads weighed 3.8 pounds, with the lowest marketable yield observed in plants spaced at 9 inches and receiving low fertilizer rates [69].

A research designed to carry out a two-season trial on two cabbage (*Brassica oleracea* var. *capitata* L.) cultivars in spring and autumn in the Beijing-Tianjin-Hebei region (2017) stated that the application of drip irrigation under

mulch should be approximately 114.7–125.0 mm and the N fertilization should be about 200 kg/ha [70]. In addition, a study conducted to determine the effect of fertilizer type on the growth and yield of two cabbage varieties (Copenhagen market and F1 minor) in Nigeria (2009 and 2010) reported that compared with NPK (15:15:15), NPK enhanced the optimum yield of cabbage varieties [71]. Moreover, the Copenhagen market variety outperformed F1 minor in head yield regardless of fertilizer application, making it the preferred choice in Ogbomoso, Southwest Nigeria, with field experiments demonstrating that fertilizing cabbage plants with 80 kg N/ha alongside 100 ppm fulvic acid results in high-quality heads suitable for local consumption in both growing seasons [72].

Combining bioslurry and NP inorganic fertilizers in southern Ethiopia significantly enhanced cabbage growth, with the study recommending the application of 50 m³/ha of bioslurry alongside 75% NP for optimal production under similar conditions [73]. Furthermore, a study conducted to assess the response of head cabbage to different rates of inorganic nitrogen fertilizer and farmyard manure in Bore, Southern Oromia, Ethiopia, reported that head cabbage responded well to the combined application of N and FYM and that application of 235 kg N/ha +6 tons FYM/ha can provide the optimum cabbage head yield in the study area [13]. Nitrogen levels profoundly affect the cabbage yield by supporting crucial growth processes such as leaf and stem development, chlorophyll production, and overall plant vigor [74]. When cabbage plants receive insufficient nitrogen, they may exhibit stunted growth, reduced leaf size, and pale green foliage [75]. Excessive nitrogen can result in overly lush vegetative growth, potentially reducing yields and head size, while simultaneously increasing susceptibility to pests and diseases [76]. Optimizing cabbage yields requires testing soil for nitrogen levels and adjusting fertilizer applications accordingly to maximize growth [77]. In addition, the timing and method of nitrogen application can also influence cabbage growth and yield [78]. Split applications throughout the growing season may help maintain more consistent levels of available nitrogen for the plants [79]. The impact of nitrogen levels on cabbage yield depends on factors such as the soil type, climate, cabbage variety, and farming methods. Understanding cabbage nitrogen uptake can optimize application timing and amount [80]. Research may investigate uptake kinetics, nitrogen efficacy for sustainable practices, environmental consequences like runoff and emissions, and nutrient interactions for yield optimization [81]. In general, nitrogen is crucial for cabbage growth and yield, but proper management is a key to optimize results and prevent harm to plant health and productivity.

9. Effect of Nitrogen Levels on the Growth of Cabbage

Soil fertility, especially nitrogen levels, significantly affects cabbage production in Ethiopia, with regional variations impacting it differently, while climate conditions such as temperature and rainfall also influence nitrogen availability and uptake by cabbage plants [82]. Ethiopia has diverse climate zones, ranging from highlands to lowlands, and has an impact

on nitrogen availability [83]. The cabbage crop's growth depends on nitrogen fertilizer, with too little causing stunted growth and excess leading to lush foliage but poor root development and disease susceptibility [84]. The source of nitrogen fertilizer (e.g., urea, ammonium nitrate, and organic sources) can influence its availability to cabbage plants and the rate of nitrogen release in the soil [49]. Different cabbage varieties may have varying nitrogen requirements and responses to nitrogen levels in the soil [85]. Different cabbage varieties vary in nitrogen utilization efficiency, influenced by farming techniques such as irrigation, mulching, crop rotation, and intercropping, affecting nitrogen availability and uptake [86]. Nitrogen levels can influence the susceptibility of cabbage plants to pests and diseases [87]. High nitrogen levels can sometimes make plants more vulnerable to certain pests and diseases [88]. Environmental stresses such as drought or waterlogging can disrupt cabbage growth by affecting nitrogen uptake, with nitrogen application methods and cover crops playing vital roles in mitigating these effects [89]. Nitrogen interacts with other essential nutrients such as phosphorus, potassium, and micronutrients [90]. Imbalances in nutrient levels can affect cabbage growth and development. Therefore, comprehending and enhancing these elements could assist Ethiopian farmers in enhancing cabbage yields by efficiently controlling nitrogen levels in their fields.

10. Effect of Nitrogen Levels on Cabbage Quality

The significant and well-documented influence of nitrogen levels on cabbage quality underscores its crucial role as a nutrient essential for optimal growth and development in cabbage plants [91]. Adequate nitrogen levels result in larger, greener leaves, which contribute to the overall size and appearance of the cabbage head [92]. Cabbage plants supplied with ample nitrogen, a crucial constituent of chlorophyll vital for photosynthesis, generally yield higher, as it enhances the plant's energy production and overall yield potential [93]. The availability of nitrogen directly impacts the size and firmness of cabbage heads, with insufficient levels leading to smaller heads with loose leaves and excessive levels causing overly large but less compact heads [32]. Adequate nitrogen boosts cabbage's nutritional quality by spurring the synthesis of proteins, vitamins, and essential nutrients, enriching its overall value [94]. Optimal nitrogen levels bolster cabbage plants against pests and diseases by fostering robust growth and enhancing their ability to fend off attacks [95]. Maintaining a balanced nitrogen application is crucial to prevent delayed maturity, increased susceptibility to diseases, and environmental pollution via groundwater leaching [96]. In addition, the timing and method of nitrogen application play a crucial role in optimizing cabbage quality and minimizing potential negative impacts [42].

11. Chemical Composition of Nitrogen Fertilizer

Nitrogen fertilizers can come in various forms, each with its own chemical composition. Urea is a solid nitrogen fertilizer with a high nitrogen content, typically around 46% [97]. It is made up of carbon, nitrogen, oxygen, and hydrogen atoms

[98]. Urea, rich in nitrogen, is a cost-effective fertilizer commonly used for cabbage plants in Ethiopia [99]. Ammonium nitrate (NH_4NO_3), a crystalline salt compound, comprises nitrogen in both ammonium and nitrate forms, constituting approximately 34% of the nitrogen content [100]. Ammonium sulfate ($(\text{NH}_4)_2\text{SO}_4$) fertilizer contains both nitrogen and sulfur, and it is composed of nitrogen, hydrogen, sulfur, and oxygen atoms [101]. Anhydrous ammonia (NH_3) is also a gas, not a solid like the others, and is often applied directly to the soil [102]. In general, these are some of the main nitrogen fertilizers used in agriculture, each with its own chemical composition and properties.

12. Summary and Conclusions

Cabbage (*Brassica oleracea* var. *capitata* L.) plays a pivotal role in Ethiopian agriculture, but its productivity hinges on nuanced factors, notably nitrogen fertilizer levels. Cabbage cultivation in Ethiopia grapples with nitrogen fertilizer management challenges, balancing the need for enhanced growth and yield against the risks of adverse effects from both excessive and moderate to high levels of nitrogen application, impacting overall yield and quality. Also, integrated nutrient management approaches, combining nitrogen fertilizers with organic amendments or other nutrients, show promise in optimizing cabbage production while minimizing environmental impacts. However, there is a lack of specific studies on how nitrogen fertilizer levels affect cabbage production in Ethiopia, indicating a need for further research. Bridging the gap between research findings and practical application, particularly in optimizing cabbage production through integrated nutrient management approaches tailored to local conditions, is crucial for supporting sustainable agricultural development in Ethiopia.

13. Recommendations

Design controlled experiments to evaluate varying nitrogen fertilizer levels' effects on cabbage yield, quality, and environmental impact in Ethiopian growing conditions using randomized controlled trials with replicated plots. Utilize isotope tracing to track nitrogen uptake by cabbage plants and understand nitrogen use efficiency. Establish long-term field trials to monitor effects on soil fertility, productivity, and environmental outcomes over multiple growing seasons. Employ a multidisciplinary approach integrating agronomy, soil science, environmental science, economics, and social sciences to comprehensively address nitrogen fertilizer management complexities. Engage stakeholders, including farmers and researchers, by integrating indigenous knowledge and traditional practices to ensure relevance and participation in research. Conduct economic analyses and develop tools to demonstrate nitrogen fertilizer management strategies' viability, while promoting knowledge exchange for improved practices.

Data Availability

Data sharing is not applicable to this article as no new data were analyzed in this study.

Conflicts of Interest

The author declares that there are no conflicts of interest.

Acknowledgments

The author would like to acknowledge Anbesu Jima for being the major source to write this review.

References

- [1] G. R. Dixon, "Vegetable Brassicas and Related Crucifers: *Crop Production Science In Horticulture Series, No. 14*," 2007, https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=G.+R.+Dixon%2C+Vegetable+brassicas+and+related+crucifers%3A+2nd+Edition%2C+CABI%2C+2007&btnG=.
- [2] K. L. H. Semuli, *Nitrogen Requirements for Cabbage (Brassica Oleracea Capitata) Transplants and Crop Response to Spacing and Nitrogen Top-Dressing*, University of Pretoria, Hatfield, Pretoria, 2005.
- [3] M. Chiang, C. Chong, B. Landry, and R. Crete, "Cabbage: *Brassica oleracea* subsp. *capitata* L.," in *Genetic Improvement of Vegetable Crops*, pp. 113–155, Elsevier, Amsterdam, Netherlands, 1993.
- [4] Y. Yebirzaf, "Effect of different rate of nitrogen fertilizer on the growth and yield of cabbage (*Brassica oleracea* L.) at Debre Markos, North West Ethiopia," *African Journal of Plant Science*, vol. 11, no. 7, pp. 276–281, 2017.
- [5] D. Atnafu, "Overview of different rate of nitrogen application on growth and yield components of head cabbage (*Brassica oleracea. capitata* L.) in Ethiopia," *American Journal of Life Sciences*, vol. 8, no. 6, p. 196, 2020.
- [6] N. Moreb, A. Murphy, S. Jaiswal, and A. K. Jaiswal, "Cabbage," *Nutritional composition and antioxidant properties of fruits and vegetables*, pp. 33–54, 2020.
- [7] M. Drozdowska, T. Leszczyńska, A. Koronowicz, E. Piasna-Słupecka, D. Domagała, and B. Kusznierevicz, "Young shoots of red cabbage are a better source of selected nutrients and glucosinolates in comparison to the vegetable at full maturity," *European Food Research and Technology*, vol. 246, no. 12, pp. 2505–2515, 2020.
- [8] S. Kumar, S. Sharma, V. Kumar, R. Sharma, A. Minhas, and R. Boddu, "Cruciferous vegetables: a mine of phytonutrients for functional and nutraceutical enrichment," in *Current Advances for Development of Functional Foods Modulating Inflammation and Oxidative Stress*, pp. 401–426, Elsevier, Amsterdam, Netherlands, 2022.
- [9] A. H. Abdumannabovna and T. S. Zakirjonovna, "The role of cabbage in the prevention of tumor diseases," *Scientific Impulse*, vol. 1, pp. 680–682, 2022.
- [10] J. Červenski, S. Vlajić, M. Ignjatov, G. Tamindžić, and S. Zec, "Agroclimatic conditions for cabbage production," *Ratarstvo i povrtarstvo*, vol. 59, no. 2, pp. 43–50, 2022.
- [11] S. Paudel, L. P. Sah, M. Devkota, V. Poudyal, P. Prasad, and M. R. Reyes, "Conservation agriculture and integrated pest management practices improve yield and income while reducing labor, pests, diseases and chemical pesticide use in smallholder vegetable farms in Nepal," *Sustainability*, vol. 12, no. 16, p. 6418, 2020.
- [12] S. Teshome and A. Amide, "Response of head cabbage (*Brassica oleracea* L.) yield to application of different rates of inorganic nitrogen and phosphorus fertilizers at Bore, Southern Ethiopia," *International Journal of Biochemistry*,

- Biophysics and Molecular Biology*, vol. 5, no. 1, pp. 18–28, 2020.
- [13] S. Teshome Wolde, D. T. Tana, and D. W. Mohammed, *Response of head cabbage (Brassica oleracea L.) to different rates of nitrogen fertilizer and farmyard manure at bore, Southern Ethiopia*, Haramaya university, Dire Dawa, Ethiopia, 2015.
 - [14] C. Suruban, M. A. Kader, and Z. M. Solaiman, "Influence of various composted organic amendments and their rates of application on nitrogen mineralization and soil productivity using Chinese cabbage (*Brassica rapa* L. var. *Chinensis*) as an indicator crop," *Agriculture*, vol. 12, no. 2, p. 201, 2022.
 - [15] D. Sarkar, A. Sankar, O. S. Devika et al., "Optimizing nutrient use efficiency, productivity, energetics, and economics of red cabbage following mineral fertilization and biopriming with compatible rhizosphere microbes," *Scientific Reports*, vol. 11, no. 1, 2021.
 - [16] F. Agostini, F. Tei, M. Silgram, M. Farneselli, P. Benincasa, and M. Aller, "Decreasing nitrate leaching in vegetable crops with better N management," *Genetic engineering, bio-fertilisation, soil quality and organic farming*, pp. 147–200, 2010.
 - [17] C. Ifejika Speranza, "Resilient adaptation to climate change in African agriculture: studies," 2010, <https://hdl.handle.net/10419/199179>.
 - [18] R. Ruggles Gates, "Wild cabbages and the effects of cultivation," *Journal of Genetics*, vol. 51, no. 2, pp. 363–372, 1953.
 - [19] S. A. A. Jahn, "The traditional domestication of a multi-purpose tree *Moringa stenopetala* (Bak. f.) Cuf. in the Ethiopian Rift Valley," *Ambio*, vol. 20, pp. 244–247, 1991.
 - [20] T. F. D. Belew, "A review on production status and consumption pattern of vegetable in Ethiopia," *Journal of Biology, Agriculture and Healthcare*, vol. 5, no. 21, 2015.
 - [21] S. Teshome and T. Bobo, "Adaptability studies of head cabbage (*Brassica oleracea* L.) varieties at adola rede areas, southern Oromia, Ethiopia," *International Journal of Asian and African Studies*, vol. 51, pp. 14–19, 2018.
 - [22] B. Bati, A. Negesse, S. Gizachew, and Y. Esma'el, "Determinants of technical efficiency of head cabbage producer farmers in west arsi zone, Oromia region, Ethiopia," *American Journal of Plant Biology*, vol. 8, p. 9.
 - [23] E. J. Ryder and E. J. Ryder, "Cabbage," *Leafy Salad Vegetables*, Springer, Berlin, Germany, 1979.
 - [24] C. W. Li, "The origin, evolution, taxonomy and hybridization of Chinese cabbage," in *Proceedings of the Chinese Cabbage 1st International Symposium*, pp. 3–10, Tsukuba Science City, Japan, April 1981.
 - [25] Z. Fang, Y. Liu, P. Lou, and G. Liu, "Current trends in cabbage breeding," *Journal of New Seeds*, vol. 6, no. 2-3, pp. 75–107, 2004.
 - [26] F. Toro, L. Martin-Closas, and A. Pelacho, "Jasmonates promote cabbage (*Brassica oleracea* L. var. *Capitata* L.) root and shoot development," in *Proceedings of the Roots: The Dynamic Interface between Plants and the Earth: The 6th Symposium of the International Society of Root Research, 11–15 November 2001*, pp. 77–83, Nagoya, Japan, November 2003.
 - [27] M. Gil, J. Tudela, A. Martínez-Sánchez, and M. Luna, "Harvest maturity indicators of leafy vegetables," *Stewart Postharvest Review*, vol. 8, pp. 1–9, 2012.
 - [28] S. Asredie Kolech and F. Mengistu Tiruneh, *Techniques of Cool Season Vegetable Crops Seed Production: A Guide for Seed Producers and Development Practitioners*, 2010, https://www.researchgate.net/profile/Semagn-Kolech/publication/330926724_Techniques_of_Cool_Season_Vegetable_Crops_Seed_Production_A_Guide_for_Seed_producers_and_Development_Practitioners/links/5c5c124192851c48a9c05ec2/Techniques-of-Cool-Season-Vegetable-Crops-Seed-Production-A-Guide-for-Seed-producers-and-Development-Practitioners.pdf.
 - [29] B. Singh, S. Sharma, P. Kalia, and B. Singh, "Genetic variability for antioxidants and horticultural traits in cabbage," *Indian Journal of Horticulture*, vol. 68, pp. 51–55, 2011.
 - [30] J. Šutković, P. Glamočlija, L. Karić, and A. Yildirim, "Genetic Characterization of *Brassica oleracea* L. var. *acephala* DC cultivars from Bosnia and Herzegovina," *Genetics and Applications*, vol. 5, no. 1, pp. 40–50, 2021.
 - [31] Csa, "Agricultural sample survey 2020/2021," *Report on Area and Production of Major Crops (Private Peasant Holdings, Meher Season)*, Central Statistical Agency, Addis Ababa, Ethiopia, 2021.
 - [32] F. Terefa, *Effect of Irrigation Scheduling and Nitrogen Fertilizer Rate on Growth and Production of Cabbage (Brassica oleracea L. Var. Capitata) in Seka District, Jimma Zone, Ethiopia*, Jimma University, Jimma, Ethiopia, 2017.
 - [33] A. Constanta and C. Lenuta, "Assessment of storage capacity of some white cabbage varieties according to histological and histochemical structure," *Journal of Horticulture, Forestry and Biotechnology*, vol. 18, pp. 18–24, 2014.
 - [34] M. Suchánková, Z. Kapounova, M. Dofkova, J. Ruprich, J. Blahova, and I. Kouřilová, "Selected fruits and vegetables: comparison of nutritional value and affordability," *Czech Journal of Food Sciences*, vol. 33, no. 3, pp. 242–246, 2015.
 - [35] M. Traka and R. Mithen, "Glucosinolates, isothiocyanates and human health," *Phytochemistry Reviews*, vol. 8, no. 1, pp. 269–282, 2009.
 - [36] J. Słupski, E. Bernaś, and P. Gębczyński, "Vegetables and vegetable-based dishes in the rural tradition of the Małopolska region," *Indicators of change in cultural heritage*, vol. 20, pp. 119–132, 2021.
 - [37] A. F. Belay and B. Gashaw, "Evaluation of levels of nps fertilizer on head cabbage (*Brassica oleracea* var. *capitata* L.) production during the two main growing seasons in Chiha district, Ethiopia," *International Journal of Agricultural and Natural Sciences*, vol. 16, pp. 231–242, 2023.
 - [38] L. G. Paranhos, C. E. Barrett, L. Zotarelli, R. Darnell, K. Migliaccio, and T. Borisova, "Planting date and in-row plant spacing effects on growth and yield of cabbage under plastic mulch," *Scientia Horticulturae*, vol. 202, pp. 49–56, 2016.
 - [39] R. Srinivasan, K. J. Rao, V. Sailaja, and D. Kalaivanan, "Influence of organic manures and fertilizers on nutrient uptake, yield and quality in cabbage-baby corn cropping sequence," *Journal of Horticultural Science*, vol. 9, no. 1, pp. 48–54, 2014.
 - [40] J.-E. Lee, P. Wang, G.-Y. Kim et al., "Effects of soil pH on nutritional and functional components of Chinese cabbage (*Brassica rapa* ssp. *campestris*)," *Horticultural Science and Technology*, vol. 28, pp. 353–362, 2010.
 - [41] S. Pramanik, "Effect of nitrogen and phosphorus on the growth and yield of cabbage (*Brassica oleracea* var. *capitata* L.)," *Department of Horticulture*, vol. 3, 2007.
 - [42] Z.-m. Qu, X.-c. Qi, J. Wang, Q. Chen, and C.-l. Li, "Effects of nitrogen application rate and topdressing times on yield and

- quality of Chinese cabbage and soil nitrogen dynamics," *Environmental Pollutants and Bioavailability*, vol. 31, pp. 1–8, 2019.
- [43] T. Salo, "Effects of band placement and nitrogen rate on dry matter accumulation, yield and nitrogen uptake of cabbage, carrot and onion," *Agricultural and Food Science*, vol. 8, no. 2, pp. 157–232, 1999.
- [44] S. Teshome and A. Amide, "Response of head cabbage (*Brassica oleracea* L.) yield to different rates of inorganic nitrogen and phosphorus fertilizers at bore, southern Ethiopia," in *Proceedings of the Regional Review Workshop on Completed Research Activities*, p. 304, Adami Tulu, Ethiopia, October 2021.
- [45] L. L. Harter and L. R. Jones, *Cabbage Diseases: US Department of Agriculture*, 1928, https://books.google.com.et/books/about/Cabbage_Diseases.html?id=tPvmGX1adIC&redir_esc=y.
- [46] B. Singh, K. Pathak, K. Sarma, and M. Thapa, "Effect of transplanting dates on plant growth, yield and quality traits of cabbage (*Brassica oleracea* var. *capitata* L.) cultivars," *Indian Journal of Hill Farming*, vol. 23, pp. 1–5, 2010.
- [47] D. Cruz Nieto, L. Montes, I. Lirión Rodríguez et al., "Analysis of the biological and chemical characteristics of cabbage (*Brassica oleracea* var. *capitata* L.) Fed with compost based on sugarcane residues," *Brazilian Journal of Biology*, vol. 83, 2023.
- [48] D. Tilman and M. Clark, "Food, agriculture and the environment: can we feed the world and save the earth?" *Dædalus*, vol. 144, no. 4, pp. 8–23, 2015.
- [49] M. Turan and F. Sevimli, "Influence of different nitrogen sources and levels on ion content of cabbage (*Brassica oleracea* var. *capitata*)," *New Zealand Journal of Crop and Horticultural Science*, vol. 33, no. 3, pp. 241–249, 2005.
- [50] W. Liu, M. Muzolf-Panek, and T. Kleiber, "Effect of nitrogen nutrition and planting date on the yield and physicochemical parameters of flowering Chinese cabbage," *Agronomy*, vol. 12, no. 11, p. 2869, 2022.
- [51] A. S. Iqbal and M. Asif, "Performance of different varieties and fertilizer management practices on the growth and yield of knol-khol," *International Journal of Agricultural Sciences*, vol. 8, 2008.
- [52] H. Akand, K. Mazed, A. Pulok, J. Moonmoon, and S. Partho, "Influence of different dose of nitrogen on the growth and yield of cabbage (*Brassica oleracea* var. *capitata* L.)," *International Journal of Multidisciplinary Research and Development*, vol. 2, pp. 11–14, 2015.
- [53] M. A. Ackah, *Mineral Nitrogen and Salicylic Acid Rates and Their Effects on Yield, Quality and Alternaria Blight Incidence in Cabbage*, University of Cape Coast, Cape Coast, Ghana, 2021.
- [54] E. Kolota and P. Chohura, "Control of head size and nutritional value of cabbage by plant population and nitrogen fertilization," *Acta Scientiarum Polonorum Hortorum Cultus*, vol. 14, pp. 75–85, 2015.
- [55] N. Fageria and A. Moreira, "The role of mineral nutrition on root growth of crop plants," *Advances in Agronomy*, vol. 110, pp. 251–331, 2011.
- [56] M. S. Shahariar, M. Moniruzzaman, B. Saha, G. Chakraborty, M. Islam, and S. Tahsin, "Effects of fresh and digested cowdung and poultry litter on the growth and yield of cabbage (*Brassica oleracea*)," *Bangladesh Journal of Scientific and Industrial Research*, vol. 48, pp. 1–6, 2013.
- [57] K. E. Mwanga, *Effects of bio-slurry and farm yard manure on soil amelioration and Chinese cabbage (Brassica Rapa Var. Chinensis) yields in Njombe region, Tanzania*, Sokoine University of Agriculture, Njombe, Tanzania, 2016.
- [58] M. D. Kleinhenz and A. Wszelaki, "Yield and relationships among head traits in cabbage as influenced by planting date and cultivar. I. Fresh market," *HortScience*, vol. 38, no. 7, pp. 1349–1354, 2003.
- [59] S. Chaudhary, S. Yadav, D. Mahto, R. Sharma, and M. Kumar, "Response of growth, yield attributes and yield of cabbage (*Brassica oleracea* var. *capitata*) to different organic and inorganic sources of nutrients in Magadha plain of Bihar," *International Journal of Current Microbiology and Applied Sciences*, pp. 4748–4756, 2018.
- [60] E. Atanasova, I. Mitova, I. Dimitrov, and I. Stancheva, "Effect of different fertilizer sources on the quality of head cabbage," *Journal of Applied Horticulture*, vol. 09, no. 01, pp. 74–76, 2007.
- [61] B. Durenne, A. Blondel, N. Ducat, O. Pigeon, M.-L. Fauconnier, and P. Druart, "Oral communication: Brassica napus L. plantlets affected during in vitro growth by the presence of epoxiconazole," *Brassica*, vol. 2017, 2017.
- [62] N. Hasan, *Efficacy of Vermicompost and Biochar on the Growth and Yield of Green Cabbage*, Department Of Horticulture, Sher-E-Bangla Agricultural University, Dhaka, Bangladesh, 2017.
- [63] P. Y. Debrah, C. Afriyie-Debrah, F. Nuhu, P. Ribeiro, and E. A. Obeng, "Response of cabbage (*Brassica oleracea* var. *capitata* L.) to organic and inorganic fertilizer," *Journal of Experimental Agriculture International*, vol. 43, pp. 105–114, 2021.
- [64] S. A. Bardisi and M. A. Mandour, "Growth and productivity of two cabbage cultivars grown in sandy soil as affected by foliar spray with npk fertilizer and amino acids," *Journal of Plant Production*, vol. 11, pp. 1063–1069, 2020.
- [65] F. A. Parry, W. A. Dar, B. A. Bhat, and T. Sultan, "Effect of different combinations of Farm Yard Manure and Bio-fertilizers on growth, yield and quality of Chinese cabbage under high altitude climatic conditions of Gurez-J&K," *International journal of chemical studies*, vol. 9, no. 1, pp. 856–858, 2021.
- [66] M. Ferré Gras, "Desarrollo de una web segura," 2023, <http://hdl.handle.net/2445/200588>.
- [67] M. S. Reza, A. Islam, M. A. Rahman, M. Y. Miah, S. Akhter, and M. M. Rahman, "Impact of organic fertilizers on yield and nutrient uptake of cabbage (*Brassica oleracea* var. *capitata*)," *Journal of Science, Technology and Environment Informatics*, vol. 3, no. 2, pp. 231–244, 2016.
- [68] A. N. Mbatha, *Influence of Organic Fertilisers on the Yield and Quality of Cabbage and Carrots*, University of the Free State, Bloemfontein, South Africa, 2008.
- [69] J. White and R. Forbes, "Effect of spacing and fertilizer rates on cabbage yield and head weight," in *Proceedings of the Florida State Horticultural Society*, pp. 118–119, Florida, FL, USA, April 1976.
- [70] X. Wu, M. Bai, Y. Li et al., "The effect of fertigation on cabbage (*Brassica oleracea* L. var. *Capitata*) grown in a greenhouse," *Water*, vol. 12, no. 4, p. 1076, 2020.
- [71] J. Olaniyi and A. Ojetayo, "Effect of fertilizer types on the growth and yield of two cabbage varieties," *Journal of Animal and Plant Sciences*, vol. 12, pp. 1573–1582, 2011.
- [72] A. Mahmoud, A. R. Abdel- Razik, S. M. El-Araby, and S. M. Ragab, "Effect of nitrogen fertilizer and fulvic acid application on the growth, productivity and nutritional quality of cabbage," *Alexandria Science Exchange Journal*, vol. 42, no. 4, pp. 921–933, 2021.

- [73] T. T. Kulelo, "Agronomic, yield and quality characteristics of cabbage (*Brassica oleracea* L. var. *Capitata*) as affected by combined application of organic and inorganic fertilizers at hawassa, southern Ethiopia," 2017, https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=T.+T.+KULELO%2C+%22Agronomic%2C+Yield+and+Quality+Characteristics+of+Cabbage+%28Brassica+oleracea+L.+var.+capitata%29+as+Affected+by+Combined+Application+of+Organic+and+Inorganic+Fertilizers+at+Hawassa%2C.
- [74] M. Reich, A. Van Den Meerakker, S. Parmar, M. Hawkesford, and L. De Kok, "Temperature determines size and direction of effects of elevated CO₂ and nitrogen form on yield quantity and quality of Chinese cabbage," *Plant Biology*, vol. 18, no. S1, pp. 63–75, 2016.
- [75] A. Alemayehu and W. Bewket, "Local climate variability and crop production in the central highlands of Ethiopia," *Environmental Development*, vol. 19, pp. 36–48, 2016.
- [76] V. Torres-Olivar, O. G. Villegas-Torres, M. L. Domínguez-Patiño et al., "Role of nitrogen and nutrients in crop nutrition," *Journal of Agricultural Science and Technology A*, vol. 4, p. 29, 2014.
- [77] H. Riley and G. Guttormsen, "N requirements of cabbage crops grown on contrasting soils. I. Field trials," *Norwegian Journal of Agricultural Sciences*, vol. 7, p. 275, 1993.
- [78] L. Brandsæter, J. Netland, and R. Meadow, "Yields, weeds, pests and soil nitrogen in a white cabbage-living mulch system," *Biological Agriculture and Horticulture*, vol. 16, no. 3, pp. 291–309, 1998.
- [79] S. Westerveld, M. McDonald, C. Scott-Dupree, and A. McKeown, "The effect of nitrogen on insect and disease pests of onions, carrots, and cabbage," *Journal of Vegetable Crop Production*, vol. 8, pp. 87–99, 2003.
- [80] D. Huber, V. Römhild, and M. Weinmann, "Relationship between nutrition, plant diseases and pests," in *Marschner's mineral Nutrition of Higher Plants*, pp. 283–298, Elsevier, Amsterdam, Netherlands, 2012.
- [81] K.-U. Katroschan, R. Uptmoor, and H. Stützel, "Nitrogen use efficiency of organically fertilized white cabbage and residual effects on subsequent beetroot," *Plant and Soil*, vol. 382, no. 1–2, pp. 237–251, 2014.
- [82] U. R. Malvi, "Interaction of micronutrients with major nutrients with special reference to potassium," *Karnataka Journal of Agricultural Sciences*, vol. 24, 2011.
- [83] K. Karthika, P. S. Philip, and S. Neenu, "Brassicaceae plants response and tolerance to nutrient deficiencies," *The Plant Family Brassicaceae: Biology and Physiological Responses to Environmental Stresses*, 2020.
- [84] S. Abbas, M. T. Javed, Q. Ali, M. Azeem, and S. Ali, "Nutrient deficiency stress and relation with plant growth and development," in *Engineering Tolerance in Crop Plants against Abiotic Stress*, pp. 239–262, CRC Press, Boca Raton, FL, USA, 2021.
- [85] T. Winch, *Growing Food: A Guide to Food Production*, Springer Science and Business Media, Berlin, Germany, 2007.
- [86] H. W. Olf, K. Blankenau, F. Brentrup, J. Jasper, A. Link, and J. Lammel, "Soil- and plant-based nitrogen-fertilizer recommendations in arable farming," *Journal of Plant Nutrition and Soil Science*, vol. 168, no. 4, pp. 414–431, 2005.
- [87] T. Coolong, K. Cassity-Duffey, and A. L. B. Ribeiro da Silva, "Influence of nitrogen rate, fertilizer type, and application method on cabbage yield and nutrient concentrations," *HortTechnology*, vol. 32, no. 2, pp. 134–139, 2022.
- [88] L. López-Bellido, R. J. López-Bellido, and R. Redondo, "Nitrogen efficiency in wheat under rainfed Mediterranean conditions as affected by split nitrogen application," *Field Crops Research*, vol. 94, no. 1, pp. 86–97, 2005.
- [89] L. Van Eerd, J. Turnbull, C. Bakker, R. Vyn, A. McKeown, and S. Westerveld, "Comparing soluble to controlled-release nitrogen fertilizers: storage cabbage yield, profit margins, and N use efficiency," *Canadian Journal of Plant Science*, vol. 98, no. 4, pp. 815–829, 2018.
- [90] H. Kubota, M. Iqbal, S. Quideau, M. Dyck, and D. Spaner, "Agronomic and physiological aspects of nitrogen use efficiency in conventional and organic cereal-based production systems," *Renewable Agriculture and Food Systems*, vol. 33, no. 5, pp. 443–466, 2018.
- [91] J. Song, J. Yang, and B. R. Jeong, "Growth, quality, and nitrogen assimilation in response to high ammonium or nitrate supply in cabbage (*Brassica campestris* L.) and lettuce (*Lactuca sativa* L.)," *Agronomy*, vol. 11, no. 12, p. 2556, 2021.
- [92] Y. Amagai, N. Lu, E. Hayashi et al., "External green light as a new tool to change colors and nutritional components of inner leaves of head cabbages," *Journal of Food Measurement and Characterization*, vol. 16, pp. 269–280, 2022.
- [93] N. S. Tehulie and S. Belete, "Review on the effects of NPS fertilizer rates on growth, yield components and yield of cabbage (*Brassica oleracea* var. *capitata* L.)," *South Asia: Journal of South Asian Studies*, vol. 1, no. 1, pp. 15–21, 2021.
- [94] M. Haghighi, A. Barzegar Sadeghabad, and R. Abolghasemi, "Effect of exogenous amino acids application on the biochemical, antioxidant, and nutritional value of some leafy cabbage cultivars," *Scientific Reports*, vol. 12, no. 1, 2022.
- [95] F. M. Bradley, *Rodale's Vegetable Garden Problem Solver: The Best and Latest Advice for Beating Pests, Diseases, and Weeds and Staying a Step Ahead of Trouble in the Garden*: Rodale Books, 2007.
- [96] M. Anas, F. Liao, K. K. Verma et al., "Fate of nitrogen in agriculture and environment: agronomic, eco-physiological and molecular approaches to improve nitrogen use efficiency," *Biological Research*, vol. 53, pp. 47–20, 2020.
- [97] B. Beig, M. B. K. Niazi, Z. Jahan, A. Hussain, M. H. Zia, and M. T. Mehran, "Coating materials for slow release of nitrogen from urea fertilizer: a review," *Journal of Plant Nutrition*, vol. 43, no. 10, pp. 1510–1533, 2020.
- [98] P. Vaughan and J. Donohue, "The structure of urea. Interatomic distances and resonance in urea and related compounds," *Acta Crystallographica*, vol. 5, no. 4, pp. 530–535, 1952.
- [99] T. Terefe, T. Ayalew, and H. Mohammed B, "Combined application of bioslurry and inorganic fertilizers on quality traits of cabbage and soil properties," *Asian Journal of Biological Sciences*, vol. 11, no. 1, pp. 24–32, 2017.
- [100] J. C. Oxley, J. L. Smith, E. Rogers, and M. Yu, "Ammonium nitrate: thermal stability and explosivity modifiers," *Thermochimica Acta*, vol. 384, no. 1–2, pp. 23–45, 2002.
- [101] L. R. Westbrook, "Process of Making Ammonium Sulphate," 1935, <https://patentimages.storage.googleapis.com/42/9a/63/1914b49bad41a9/US1989124.pdf>.
- [102] D. L. Mangum, "Influence of anhydrous ammonia on the solubility of soil organic matter," 1965, https://ir.library.oregonstate.edu/concern/graduate_thesis_or_dissertations/vt150n35b.

- [103] J. Andoloro, K. Rose, A. Shelton, C. Hoy, and R. Becker, "Cabbage growth stages," 1983, <https://ecommons.cornell.edu/server/api/core/bitstreams/63b76faa-da7e-469d-b4fb-f58f1f6bd983/content>.
- [104] C. North, "Studies in Morphogenesis of *Brassica oleracea* L: I. Growth and development of cabbage during the vegetative phase," *Journal of Experimental Botany*, vol. 8, no. 2, pp. 304–312, 1957.
- [105] T. Hara and Y. Sonoda, "The role of macronutrients for cabbage-head formation (preliminary report) Growth performance of a cabbage plant, and potassium nutrition in the plant," *Soil Science and Plant Nutrition*, vol. 25, no. 1, pp. 103–111, 1979.